

Resources and References Handout

Medication Management in Older Adults

Hyperlinks

<http://medstopper.com/> Accessed December 12, 2016

<http://www.adultmededucation.com/index.html> Accessed December 12, 2016

<https://www.drugs.com/mednotes/> Accessed December 12, 2016

References

American College of Gastroenterology Chronic Constipation Task Force. (2005). An evidence-based approach to the management of chronic constipation in North America. *Am J Gastroenterol*, 100 Suppl 1, S1-4. doi:10.1111/j.1572-0241.2005.50613_1.x

American Diabetes Association. (2016). 10. Older Adults. *Diabetes Care*, 39 Suppl 1, S81-85. doi:10.2337/dc16-S013

American Geriatrics Society. (2015). American Geriatrics Society 2015 Updated Beers Criteria for Potentially Inappropriate Medication Use in Older Adults. *J Am Geriatr Soc*, 63(11), 2227-2246. doi:10.1111/jgs.13702

American Pharmacists Association (Producer). (2014a). Module 3: Getting ready for MTM service delivery: knowledge and skills. *APhA delivering medication therapy management services, a national certificate training program*.

American Pharmacists Association (Producer). (2014b). Module 4: Ready for action: conducting an MTM encounter. *APhA delivering medication therapy management services, a national certificate training program*.

American Pharmacists Association and National Association of Chain Drug Stores Foundation. (2008). Medication Therapy Management in Pharmacy Practice: Core Elements of an MTM Service Model. Retrieved from http://www.pharmacist.com/sites/default/files/files/core_elements_of_an_mt_m_practice.pdf. Accessed September 13, 2016

Bain KT, Holmes HM, Beers MH, Maio V, Handler SM, & Pauker SG. (2008). Discontinuing medications: a novel approach for revising the prescribing stage of the medication-use process. *J Am Geriatr Soc*, 56(10), 1946-1952. doi:10.1111/j.1532-5415.2008.01916.x

Bedell SE, Jabbour S, Goldberg R, Glaser H, Gobble S, Young-Xu Y, Graboys TB, & Ravid S. (2000). Discrepancies in the use of medications: their extent and predictors in an outpatient practice. *Arch Intern Med*, 160(14), 2129-2134.

Bluml BM. (2005). Definition of medication therapy management: development of professionwide consensus. *J Am Pharm Assoc (2003)*, 45(5), 566-572.

Boockvar K, Fishman E, Kyriacou CK, Monias A, Gavi S, & Cortes T. (2004). Adverse events due to discontinuations in drug use and dose changes in patients transferred between acute and long-term care facilities. *Arch Intern Med*, 164(5), 545-550. doi:10.1001/archinte.164.5.545

Chrischilles EA, Carter BL, Lund BC, Rubenstein LM, Chen-Hardee SS, Voelker MD, Park TR, & Kuehl AK. (2004). Evaluation of the Iowa Medicaid pharmaceutical case management program. *J Am Pharm Assoc (2003)*, 44(3), 337-349.

Christensen M, & Lundh A. (2016). Medication review in hospitalised patients to reduce morbidity and mortality. *Cochrane Database Syst Rev*, 2, Cdo08986. doi:10.1002/14651858.CD008986.pub3

Cipolle R, Strand L, & Morley P. (1998). *Pharmaceutical Care Practice*. New York, NY: McGraw-Hill.

Claxton AJ, Cramer J, & Pierce C. (2001). A systematic review of the associations between dose regimens and medication compliance. *Clin Ther*, 23(8), 1296-1310.

Coleman EA, Smith JD, Raha D, & Min SJ. (2005). Posthospital medication discrepancies: prevalence and contributing factors. *Arch Intern Med*, 165(16), 1842-1847. doi:10.1001/archinte.165.16.1842

Cosman F, de Beur SJ, LeBoff MS, Lewiecki EM, Tanner B, Randall S, & Lindsay R. (2014). Clinician's Guide to Prevention and Treatment of Osteoporosis. *Osteoporos Int*, 25(10), 2359-2381. doi:10.1007/s00198-014-2794-2

Cowper PA, Weinberger M, Hanlon JT, Landsman PB, Samsa GP, Uttech KM, Schmader KE, Lewis IK, Cohen HJ, & Feussner JR. (1998). The cost-effectiveness of a clinical pharmacist intervention among elderly outpatients. *Pharmacotherapy*, 18(2), 327-332.

Dobrzanski S, Hammond I, Khan G, & Holdsworth H. (2002). The nature of hospital prescribing errors. *British Journal of Clinical Governance*, 7(3), 187-193. doi:doi:10.1108/14664100210438271

Dolder CR, Nelson MH, & Iler CA. (2012). The effects of mirtazapine on sleep in patients with major depressive disorder. *Ann Clin Psychiatry*, 24(3), 215-224.

Eischens KP, Gilling SW, Okerlund RE, Grund TR, Iverson PS, & Schommer JC. (2010). Improving medication therapy management through collaborative hospice care in rural Minnesota. *J Am Pharm Assoc (2003)*, 50(3), 379-383. doi:10.1331/JAPhA.2010.09043

Ellis W, Kaasalainen S, Baxter P, & Ploeg J. (2012). Medication management for nurses working in long-term care. *Can J Nurs Res*, 44(3), 128-149.

Fawcett J, & Barkin RL. (1998). Review of the results from clinical studies on the efficacy, safety and tolerability of mirtazapine for the treatment of patients with major depression. *J Affect Disord*, 51(3), 267-285.

Gurwitz J. (1995). Investigating polypharmacy presents numerous opportunities for problem solving. *Brown University Long-Term Care Quality Letter*, May 15.

Hanlon JT, Weinberger M, Samsa GP, Schmader KE, Uttech KM, Lewis IK, Cowper PA, Landsman PB, Cohen HJ, & Feussner JR. (1996). A randomized, controlled trial of a clinical pharmacist intervention to improve inappropriate

prescribing in elderly outpatients with polypharmacy. *Am J Med*, 100(4), 428-437. doi:10.1016/s0002-9343(97)89519-8

Isetts BJ, Schondelmeyer SW, Artz MB, Lenarz LA, Heaton AH, Wadd WB, Brown LM, & Cipolle RJ. (2008). Clinical and economic outcomes of medication therapy management services: the Minnesota experience. *J Am Pharm Assoc (2003)*, 48(2), 203-211; 203 p following 211. doi:10.1331/JAPhA.2008.07108

James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, Lackland DT, LeFevre ML, MacKenzie TD, Ogedegbe O, Smith SC, Jr., Svetkey LP, Taler SJ, Townsend RR, Wright JT, Jr., Narva AS, & Ortiz E. (2014). 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA*, 311(5), 507-520. doi:10.1001/jama.2013.284427

Krska J, Cromarty JA, Arris F, Jamieson D, Hansford D, Duffus PR, Downie G, & Seymour DG. (2001). Pharmacist-led medication review in patients over 65: a randomized, controlled trial in primary care. *Age Ageing*, 30(3), 205-211.

Krska J, Gill D, & Hansford D. (2006). Pharmacist-supported medication review training for general practitioners: feasibility and acceptability. *Med Educ*, 40(12), 1217-1225. doi:10.1111/j.1365-2929.2006.02633.x

Lee SS, Schwemm AK, Reist J, Cantrell M, Andreski M, Doucette WR, Chrischilles EA, & Farris KB. (2009). Pharmacists' and pharmacy students' ability to identify drug-related problems using TIMER (Tool to Improve Medications in the Elderly via Review). *Am J Pharm Educ*, 73(3), 52.

Lenaghan E, Holland R, & Brooks A. (2007). Home-based medication review in a high risk elderly population in primary care--the POLYMED randomised controlled trial. *Age Ageing*, 36(3), 292-297. doi:10.1093/ageing/afm036

Lowe CJ, Petty DR, Zermansky AG, & Raynor DK. (2000). Development of a method for clinical medication review by a pharmacist in general practice. *Pharm World Sci*, 22(4), 121-126.

Medicare prescription drug, improvement, and modernization (MMA) act of 2003, Pub. L. No. 108-173, 2066 Stat. (2003 December 8, 2003).

Morisky DE, Green LW, & Levine DM. (1986). Concurrent and predictive validity of a self-reported measure of medication adherence. *Med Care*, 24(1), 67-74.

Nieuwlaat R, Wilczynski N, Navarro T, Hobson N, Jeffery R, Keepanasseril A, Agoritsas T, Mistry N, Iorio A, Jack S, Sivaramalingam B, Iserman E, Mustafa RA, Jedraszewski D, Cotoi C, & Haynes RB. (2014). Interventions for enhancing medication adherence. *Cochrane Database Syst Rev*(11), Cd000011. doi:10.1002/14651858.CD000011.pub4

O'Mahony D, O'Sullivan D, Byrne S, O'Connor MN, Ryan C, & Gallagher P. (2015). STOPP/START criteria for potentially inappropriate prescribing in older people: version 2. *Age Ageing*, 44(2), 213-218. doi:10.1093/ageing/afu145

Patient-Centered Primary Care Collaborative. (2012). The patient-centered medical home: integrating comprehensive medication management to optimize patient outcomes. Resource guide, 2nd ed. Retrieved from <https://www.pcpcc.org/sites/default/files/media/medmanagement.pdf>. Accessed July 14, 2016

Pellegrino AN, Martin MT, Tilton JJ, & Touchette DR. (2009). Medication therapy management services: definitions and outcomes. *Drugs*, 69(4), 393-406. doi:10.2165/00003495-200969040-00001

Pfizer. (2012). Meclizine [Package Insert]. New York, NY: Author.

Qaseem A, Kansagara D, Forcica MA, Cooke M, & Denberg TD. (2016). Management of Chronic Insomnia Disorder in Adults: A Clinical Practice Guideline From the American College of Physicians. *Ann Intern Med*, 165(2), 125-133. doi:10.7326/m15-2175

Reidt S, Holtan H, Stender J, Salvatore T, & Thompson B. (2016). Integrating home-based medication therapy management (MTM) services in a health system. *J Am Pharm Assoc (2003)*, 56(2), 178-183. doi:10.1016/j.japh.2016.01.003

Schmader KE, Hanlon JT, Pieper CF, Sloane R, Ruby CM, Twersky J, Francis SD, Branch LG, Lindblad CI, Artz M, Weinberger M, Feussner JR, & Cohen HJ. (2004). Effects of geriatric evaluation and management on adverse drug reactions and suboptimal prescribing in the frail elderly. *Am J Med*, 116(6), 394-401. doi:10.1016/j.amjmed.2003.10.031

Shrank WH, Polinski JM, & Avorn J. (2007). Quality indicators for medication use in vulnerable elders. *J Am Geriatr Soc*, 55 Suppl 2, S373-382. doi:10.1111/j.1532-5415.2007.01345.x

Smith JD, Coleman EA, & Min SJ. (2004). A new tool for identifying discrepancies in postacute medications for community-dwelling older adults. *Am J Geriatr Pharmacother*, 2(2), 141-147.

Smith M, Giuliano MR, & Starkowski MP. (2011). In Connecticut: improving patient medication management in primary care. *Health Aff (Millwood)*, 30(4), 646-654. doi:10.1377/hlthaff.2011.0002

Soumerai SB, Pierre-Jacques M, Zhang F, Ross-Degnan D, Adams AS, Gurwitz J, Adler G, & Safran DG. (2006). Cost-related medication nonadherence among elderly and disabled medicare beneficiaries: a national survey 1 year before the medicare drug benefit. *Arch Intern Med*, 166(17), 1829-1835. doi:10.1001/archinte.166.17.1829

Steinman MA, & Hanlon JT. (2010). Managing medications in clinically complex elders: "There's got to be a happy medium". *JAMA*, 304(14), 1592-1601. doi:10.1001/jama.2010.1482

Steurbaut S, Leemans L, Leysen T, De Baere E, Cornu P, Mets T, & Dupont AG. (2010). Medication history reconciliation by clinical pharmacists in elderly inpatients admitted from home or a nursing home. *Ann Pharmacother*, 44(10), 1596-1603. doi:10.1345/aph.1P192

Summary of the executive sessions on medication therapy management programs. Bethesda, Maryland, June 14 and August 18, 2004. (2005). *J Manag Care Pharm*, 11(2), 179-186.

Tarumi Y, Wilson MP, Szafran O, & Spooner GR. (2013). Randomized, double-blind, placebo-controlled trial of oral docusate in the management of

constipation in hospice patients. *J Pain Symptom Manage*, 45(1), 2-13.
doi:10.1016/j.jpainsymman.2012.02.008

Tjia J, Talebreza S, Reblin M, Beck A, & Ellington L. (2014). Scramble or script: responding to new Medicare billing for medications in hospice. *J Palliat Med*, 17(10), 1085-1086. doi:10.1089/jpm.2014.0230

Tomechko MA, Strand LM, Morley PC, & Cipolle RJ. (1995). Q and A from the Pharmaceutical Care Project in Minnesota. *Am Pharm, Ns*35(4), 30-39.

U.S. Food & Drug Administration. (2011). FDA Drug Safety Communication: Possible increased risk of fractures of the hip, wrist, and spine with the use of proton pump inhibitors. Retrieved from <http://www.fda.gov/Drugs/DrugSafety/PostmarketDrugSafetyInformationforPatientsandProviders/ucm213206.htm>. Accessed September 13, 2016

U.S. Food & Drug Administration. (2015). FDA Drug Safety Communication: FDA strengthens warning that non-aspirin nonsteroidal anti-inflammatory drugs (NSAIDs) can cause heart attacks or strokes. Retrieved from <http://www.fda.gov/Drugs/DrugSafety/ucm451800.htm>. Accessed September 13, 2016

Watts NB, Bilezikian JP, Camacho PM, Greenspan SL, Harris ST, Hodgson SF, Kleerekoper M, Luckey MM, McClung MR, Pollack RP, & Petak SM. (2010). American Association of Clinical Endocrinologists Medical Guidelines for Clinical Practice for the diagnosis and treatment of postmenopausal osteoporosis. *Endocr Pract*, 16 Suppl 3, 1-37.

Weber MA, Schiffrin EL, White WB, Mann S, Lindholm LH, Kenerson JG, Flack JM, Carter BL, Materson BJ, Ram CV, Cohen DL, Cadet JC, Jean-Charles RR, Taler S, Kountz D, Townsend RR, Chalmers J, Ramirez AJ, Bakris GL, Wang J, Schutte AE, Bisognano JD, Touyz RM, Sica D, & Harrap SB. (2014). Clinical practice guidelines for the management of hypertension in the community: a statement by the American Society of Hypertension and the International Society of Hypertension. *J Clin Hypertens (Greenwich)*, 16(1), 14-26.
doi:10.1111/jch.12237

Weber V, White A, & McIlvried R. (2008). An electronic medical record (EMR)-based intervention to reduce polypharmacy and falls in an ambulatory rural elderly population. *J Gen Intern Med*, 23(4), 399-404. doi:10.1007/s11606-007-0482-z

Wiggins BS, Rodgers JE, DiDomenico RJ, Cook AM, & Page RL, 2nd. (2013). Discharge counseling for patients with heart failure or myocardial infarction: a best practices model developed by members of the American College of Clinical Pharmacy's Cardiology Practice and Research Network based on the Hospital to Home (H2H) Initiative. *Pharmacotherapy*, 33(5), 558-580. doi:10.1002/phar.1231

Williams ME, Pulliam CC, Hunter R, Johnson TM, Owens JE, Kincaid J, Porter C, & Koch G. (2004). The short-term effect of interdisciplinary medication review on function and cost in ambulatory elderly people. *J Am Geriatr Soc*, 52(1), 93-98.