

## Resources and References Handout

### **Patient-centered Dementia Care: Understanding Patient and Caregiver Experiences**

#### Dementia

Including Mild and Major Neurocognitive Disorders (NCD)

#### Hyperlinks

[https://www.alz.org/downloads/facts\\_figures\\_2013.pdf](https://www.alz.org/downloads/facts_figures_2013.pdf)

Accessed May 2, 2017

<http://www.cstdementia.com/page/guiding-principles>

Accessed May 2, 2017

<http://www.healthinaging.org/news/tip-sheets/>

Accessed May 2, 2017

#### References

Abraha I, Rimland JM, Lozano-Montoya I, Dell'Aquila G, Velez-Diaz-Pallares M, Trotta FM, Cruz-Jentoft AJ, & Cherubini A. (2017). Simulated presence therapy for dementia. *Cochrane Database Syst Rev*, 4, Cd011882.

doi:10.1002/14651858.CD011882.pub2

Allen DN. (2011). Brief cognitive rating scale. In J. S. Kreutzer, J. DeLuca, & B. Caplan (Eds.), *Encyclopedia of clinical neuropsychology* (pp. 446-447). New York, NY: Springer.

Alzheimer's Australia. (2015). Diagnostic criteria for dementia. Retrieved from [https://www.fightdementia.org.au/files/helpsheets/Helpsheet-DementiaQandA11-DiagnosticCriteriaForDementia\\_english.pdf](https://www.fightdementia.org.au/files/helpsheets/Helpsheet-DementiaQandA11-DiagnosticCriteriaForDementia_english.pdf). Accessed May 18, 2017

Alzheimer's Association. (2015). 2015 Alzheimer's disease facts and figures. *Alzheimer's Dement*, 11(3), 332-384. Available at [http://www.alz.org/facts/downloads/facts\\_figures\\_2015.pdf](http://www.alz.org/facts/downloads/facts_figures_2015.pdf). Accessed September 13, 2016.

American Music Therapy Association. (2014). Music therapy and the film, "Alive Inside". Retrieved from [http://www.musictherapy.org/music\\_therapy\\_and\\_the\\_film\\_alive\\_inside/](http://www.musictherapy.org/music_therapy_and_the_film_alive_inside/). Accessed May 18, 2017

American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Washington, DC: Author.

American Psychiatric Association. (2014). Guideline watch (October 2014): practice guideline for the treatment of patients with Alzheimer's disease and other dementias. Retrieved from [http://psychiatryonline.org/pb/assets/raw/sitewide/practice\\_guidelines/guidelines/alzheimerwatch.pdf](http://psychiatryonline.org/pb/assets/raw/sitewide/practice_guidelines/guidelines/alzheimerwatch.pdf). Accessed May 18, 2017

American Psychiatric Association. (2007). Practice guideline for the treatment of patients with Alzheimer's disease and other dementias: second edition. Retrieved from

[http://psychiatryonline.org/pb/assets/raw/sitewide/practice\\_guidelines/guidelines/alzheimers.pdf](http://psychiatryonline.org/pb/assets/raw/sitewide/practice_guidelines/guidelines/alzheimers.pdf). Accessed May 18, 2017

Anjum A. (2016). The amnesic syndrome. In S. H. Fatemi & P. J. Clayton (Eds.), *The medical basis of psychiatry* (pp. 43-51). New York, NY: Springer.

Aripiprazole [package insert]. (2016). Carmel, NY: Silarx Pharmaceuticals, Inc.

Auchus AP, Brashear HR, Salloway S, Korczyn AD, De Deyn PP, & Gassmann-Mayer C. (2007). Galantamine treatment of vascular dementia: a randomized trial. *Neurology*, *69*(5), 448-458. doi:10.1212/01.wnl.0000266625.31615.f6

Bahar-Fuchs A, Clare L, & Woods B. (2013). Cognitive training and cognitive rehabilitation for persons with mild to moderate dementia of the Alzheimer's or vascular type: a review. *Alzheimer's Res Ther*, *5*(4), 35. doi:10.1186/alzrt189

Ballard C, Sauter M, Scheltens P, He Y, Barkhof F, van Straaten EC, van der Flier WM, Hsu C, Wu S, & Lane R. (2008). Efficacy, safety and tolerability of rivastigmine capsules in patients with probable vascular dementia: the VantagE study. *Curr Med Res Opin*, *24*(9), 2561-2574. doi:10.1185/03007990802328142

Beach SR, Celano CM, Noseworthy PA, Januzzi JL, & Huffman JC. (2013). QTc prolongation, torsades de pointes, and psychotropic medications. *Psychosomatics*, *54*(1), 1-13. doi:10.1016/j.psych.2012.11.001

Blazer DG, Yaffe K, & Karlawish J. (2015). Cognitive aging: a report from the Institute of Medicine. *JAMA*, *313*(21), 2121-2122. doi:10.1001/jama.2015.4380

Bonelli RM, & Cummings JL. (2008). Frontal-subcortical dementias. *Neurologist*, *14*(2), 100-107. doi:10.1097/NRL.0b013e31815b0de2

Borson S, Scanlan J, Brush M, Vitaliano P, & Dokmak A. (2000). The mini-cog: a cognitive 'vital signs' measure for dementia screening in multi-lingual elderly. *Int J Geriatr Psychiatry*, *15*(11), 1021-1027.

Bourgeois MS, & Hickey EM. (2009). *Dementia: From diagnosis to management- a functional approach*. New York, NY: Psychology Press.

Brodaty H, Ames D, Snowdon J, Woodward M, Kirwan J, Clarnette R, Lee E, & Greenspan A. (2005). Risperidone for psychosis of Alzheimer's disease and mixed dementia: results of a double-blind, placebo-controlled trial. *Int J Geriatr Psychiatry*, *20*(12), 1153-1157. doi:10.1002/gps.1409

Brodaty H, Ames D, Snowdon J, Woodward M, Kirwan J, Clarnette R, Lee E, Lyons B, & Grossman F. (2003). A randomized placebo-controlled trial of risperidone for the treatment of aggression, agitation, and psychosis of dementia. *J Clin Psychiatry*, *64*(2), 134-143.

Cameron DE. (1941). Studies in senile nocturnal delirium. *Psychiatric Quarterly*, *15*(1), 47-53. doi:10.1007/bf01613953

Chan WC, Lam LC, Choy CN, Leung VP, Li SW, & Chiu HF. (2001). A double-blind randomised comparison of risperidone and haloperidol in the treatment of behavioural and psychological symptoms in Chinese dementia patients. *Int J Geriatr Psychiatry*, *16*(12), 1156-1162.

Chen JH, Lamberg JL, Chen YC, Kiely DK, Page JH, Person CJ, & Mitchell SL. (2006). Occurrence and treatment of suspected pneumonia in long-term care residents dying with advanced dementia. *J Am Geriatr Soc*, *54*(2), 290-295. doi:10.1111/j.1532-5415.2005.00524.x

Chenoweth L, King MT, Jeon YH, Brodaty H, Stein-Parbury J, Norman R, Haas M, & Luscombe G. (2009). Caring for Aged Dementia Care Resident Study (CADRES) of

person-centred care, dementia-care mapping, and usual care in dementia: a cluster-randomised trial. *Lancet Neurol*, 8(4), 317-325. doi:10.1016/s1474-4422(09)70045-6

Cipriani G, Vedovello M, Ulivi M, Nuti A, & Lucetti C. (2013). Repetitive and stereotypic phenomena and dementia. *Am J Alzheimer's Dis Other Demen*, 28(3), 223-227. doi:10.1177/1533317513481094

D'Agata E, & Mitchell SL. (2008). Patterns of antimicrobial use among nursing home residents with advanced dementia. *Arch Intern Med*, 168(4), 357-362. doi:10.1001/archinternmed.2007.104

Daly SV. (2016). *The effects of visual arts on expressive language in participants with dementia [thesis]*. University of South Florida. Retrieved from <http://scholarcommons.usf.edu/etd/6215>

De Deyn P, Jeste DV, Swanink R, Kostic D, Breder C, Carson WH, & Iwamoto T. (2005). Aripiprazole for the treatment of psychosis in patients with Alzheimer's disease: a randomized, placebo-controlled study. *J Clin Psychopharmacol*, 25(5), 463-467.

De Deyn P, Rabheru K, Rasmussen A, Bocksberger JP, Dautzenberg PL, Eriksson S, & Lawlor BA. (1999). A randomized trial of risperidone, placebo, and haloperidol for behavioral symptoms of dementia. *Neurology*, 53(5), 946-955.

Deberdt WG, Dysken MW, Rappaport SA, Feldman PD, Young CA, Hay DP, Lehman DL, Dossenbach M, Degenhardt EK, & Breier A. (2005). Comparison of olanzapine and risperidone in the treatment of psychosis and associated behavioral disturbances in patients with dementia. *Am J Geriatr Psychiatry*, 13(8), 722-730. doi:10.1176/appi.ajgp.13.8.722

DiZazzo-Miller R, Pociask FD, & Samuel P. (2013). Understanding resource needs of persons with dementia and their caregivers. *Michigan Family Review*, 17(1), 1-20.

Dubois B, Padovani A, Scheltens P, Rossi A, & Dell'Agnello G. (2016). Timely diagnosis for Alzheimer's disease: a literature review on benefits and challenges. *J Alzheimer's Dis*, 49(3), 617-631. doi:10.3233/jad-150692

Fabiszewski KJ, Volicer B, & Volicer L. (1990). Effect of antibiotic treatment on outcome of fevers in institutionalized Alzheimer patients. *JAMA*, 263(23), 3168-3172.

Farlow MR, Grossberg GT, Sadowsky CH, Meng X, & Somogyi M. (2013). A 24-week, randomized, controlled trial of rivastigmine patch 13.3 mg/24 h versus 4.6 mg/24 h in severe Alzheimer's dementia. *CNS Neurosci Ther*, 19(10), 745-752. doi:10.1111/cns.12158

Farlow MR, Salloway S, Tariot PN, Yardley J, Moline ML, Wang Q, Brand-Schieber E, Zou H, Hsu T, & Satlin A. (2010). Effectiveness and tolerability of high-dose (23 mg/d) versus standard-dose (10 mg/d) donepezil in moderate to severe Alzheimer's disease: A 24-week, randomized, double-blind study. *Clin Ther*, 32(7), 1234-1251. doi:10.1016/j.clinthera.2010.06.019

Feil N. (2014). Validation therapy with late-onset dementia populations. In G. M. M. Jones & B. M. L. Misesen (Eds.), *Caregiving in dementia: Research and applications* (pp. 199-218). London: Tavistock.

Funnell E. (2000). Apperceptive agnosia and the visual recognition of object categories in dementia of the Alzheimer type. *Neurocase*, 6(6), 451-463. doi:10.1080/13554790008402716

Galvin JE, Duda JE, Kaufer DI, Lippa CF, Taylor A, & Zarit SH. (2010). Lewy body dementia: caregiver burden and unmet needs. *Alzheimer Dis Assoc Disord*, 24(2), 177-181. doi:10.1097/WAD.0b013e3181c72b5d

Gillespie P, O'Shea E, Cullinan J, Lacey L, Gallagher D, & Ni Mhaolain A. (2013). The effects of dependence and function on costs of care for Alzheimer's disease and mild cognitive impairment in Ireland. *Int J Geriatr Psychiatry*, 28(3), 256-264. doi:10.1002/gps.3819

Givens JL, Selby K, Goldfeld KS, & Mitchell SL. (2012). Hospital transfers of nursing home residents with advanced dementia. *J Am Geriatr Soc*, 60(5), 905-909. doi:10.1111/j.1532-5415.2012.03919.x

Gruss V, McCann JJ, Edelman P, & Farran CJ. (2004). Job stress among nursing home certified nursing assistants: Comparison of empowered and nonempowered work environments. *Alzheimer's Care Today*, 5(3), 207-216.

Hallenback J. (2003). *Palliative care perspectives*. New York, NY: Oxford University Press.

Harmer BJ, & Orrell M. (2008). What is meaningful activity for people with dementia living in care homes? A comparison of the views of older people with dementia, staff and family carers. *Aging Ment Health*, 12(5), 548-558. doi:10.1080/13607860802343019

Hogan DB. (2006). Donepezil for severe Alzheimer's disease. *Lancet*, 367(9516), 1031-1032. doi:10.1016/s0140-6736(06)68395-5

Huesgen A, Michel J, Rickard L, & Rickert M. (2014). Sensational Aging: Evidence on the Use of Sensory-Based Interventions for Adults with Dementia. Retrieved from

[https://www.google.com/url?sa=t&rct=j&q=&esrc=s&source=web&cd=1&ved=0ahUK EwimqIzA5vrTAhWKxlQKHdooBnwQFggpMAA&url=http%3A%2F%2Fjdc.jefferson.edu%2Fcgi%2Fviewcontent.cgi%3Ffilename%3D0%26article%3D1013%26context%3Dcreateday%26type%3Dadditional&usg=AFQjCNFN8NES9GknZGD2\\_Mfu3unK7Puafg&sig2=zxO2EGJBeAkBcYmRY3xBFA](https://www.google.com/url?sa=t&rct=j&q=&esrc=s&source=web&cd=1&ved=0ahUK EwimqIzA5vrTAhWKxlQKHdooBnwQFggpMAA&url=http%3A%2F%2Fjdc.jefferson.edu%2Fcgi%2Fviewcontent.cgi%3Ffilename%3D0%26article%3D1013%26context%3Dcreateday%26type%3Dadditional&usg=AFQjCNFN8NES9GknZGD2_Mfu3unK7Puafg&sig2=zxO2EGJBeAkBcYmRY3xBFA)

Hanyu H, Sato T, Hirao K, Kanetaka H, Sakurai H, & Iwamoto T. (2009). Differences in clinical course between dementia with Lewy bodies and Alzheimer's disease. *Eur J Neurol*, 16(2), 212-217. doi:10.1111/j.1468-1331.2008.02388.x

Hurley AC, & Volicer L. (2002). Alzheimer disease: "It's okay, Mama, if you want to go, it's okay". *JAMA*, 288(18), 2324-2331.

Kaneko Y, Butler JP, Saitoh E, Horie T, Fujii M, & Sasaki H. (2013). Efficacy of white noise therapy for dementia patients with schizophrenia. *Geriatr Gerontol Int*, 13(3), 808-810. doi:10.1111/ggi.12028

Kavirajan H, & Schneider LS. (2007). Efficacy and adverse effects of cholinesterase inhibitors and memantine in vascular dementia: a meta-analysis of randomised controlled trials. *Lancet Neurol*, 6(9), 782-792. doi:10.1016/s1474-4422(07)70195-3

Kertesz A, Morlog D, Light M, Blair M, Davidson W, Jesso S, & Brashear R. (2008). Galantamine in frontotemporal dementia and primary progressive aphasia. *Dement Geriatr Cogn Disord*, 25(2), 178-185. doi:10.1159/000113034

Khachiyants N, Trinkle D, Son SJ, & Kim KY. (2011). Sundown syndrome in persons with dementia: an update. *Psychiatry Investig*, 8(4), 275-287. doi:10.4306/pi.2011.8.4.275

Kovach CR. (2013). *End-Stage dementia care: a basic guide*. Washington, DC: Taylor & Francis.

Lach HW, & Chang YP. (2007). Caregiver perspectives on safety in home dementia care. *West J Nurs Res*, 29(8), 993-1014. doi:10.1177/0193945907303098

Lerner AJ, Hedera P, Koss E, Stuckey J, & Friedland RP. (1997). Delirium in Alzheimer disease. *Alzheimer Dis Assoc Disord*, 11(1), 16-20.

Luchins DJ, Hanrahan P, & Murphy K. (1997). Criteria for enrolling dementia patients in hospice. *J Am Geriatr Soc*, 45(9), 1054-1059.

Maglione M, Ruelaz Maher A, Hu J, Wang Z, Shanman R, Shekelle P, Roth B, Hilton L, Suttorp M, Ewing B, Motala A, & Perry T. (2011). *Off-Label use of atypical antipsychotics: an update. Comparative Effectiveness Review No. 43. (Prepared by the Southern California Evidence-based Practice Center under Contract No. HHS290-2007-10062-1)*. Retrieved from Rockville, MD:

<https://www.ncbi.nlm.nih.gov/books/NBK66081/>

Malmstrom TK, Voss VB, Cruz-Oliver DM, Cummings-Vaughn LA, Tumosa N, Grossberg GT, & Morley JE. (2015). The Rapid Cognitive Screen (RCS): a point-of-care screening for dementia and mild cognitive impairment. *J Nutr Health Aging*, 19(7), 741-744. doi:10.1007/s12603-015-0564-2

Maust DT, Kim HM, Seyfried LS, Chiang C, Kavanagh J, Schneider LS, & Kales HC. (2015). Antipsychotics, other psychotropics, and the risk of death in patients with dementia: number needed to harm. *JAMA Psychiatry*, 72(5), 438-445. doi:10.1001/jamapsychiatry.2014.3018

McKeith I, Del Ser T, Spano P, Emre M, Wesnes K, Anand R, Cicin-Sain A, Ferrara R, & Spiegel R. (2000). Efficacy of rivastigmine in dementia with Lewy bodies: a randomised, double-blind, placebo-controlled international study. *Lancet*, 356(9247), 2031-2036. doi:10.1016/s0140-6736(00)03399-7

Mitchell SL, Kiely DK, Hamel MB, Park PS, Morris JN, & Fries BE. (2004). Estimating prognosis for nursing home residents with advanced dementia. *JAMA*, 291(22), 2734-2740. doi:10.1001/jama.291.22.2734

Mitchell SL, Miller SC, Teno JM, Kiely DK, Davis RB, & Shaffer ML. (2010). Prediction of 6-month survival of nursing home residents with advanced dementia using ADEPT vs. hospice eligibility guidelines. *JAMA*, 304(17), 1929-1935. doi:10.1001/jama.2010.1572

Mitchell SL, Teno JM, Kiely DK, Shaffer ML, Jones RN, Prigerson HG, Volicer L, Givens JL, & Hamel MB. (2009). The clinical course of advanced dementia. *N Engl J Med*, 361(16), 1529-1538. doi:10.1056/NEJMoa0902234

Mitchell SL. (2007). A 93-year-old man with advanced dementia and eating problems. *JAMA*, 298(21), 2527-2536. doi:10.1001/jama.298.17.jrr70001

Mitchell SL. (2015). Advanced Dementia. *N Engl J Med*, 373(13), 1276-1277. doi:10.1056/NEJMc1509349

Mohamed S, Rosenheck R, Lyketsos CG, Kaczynski R, Sultzer DL, & Schneider LS. (2012). Effect of second-generation antipsychotics on caregiver burden in Alzheimer's disease. *J Clin Psychiatry*, 73(1), 121-128. doi:10.4088/JCP.10m06574

Mori E, Ikeda M, & Kosaka K. (2012). Donepezil for dementia with Lewy bodies: a randomized, placebo-controlled trial. *Ann Neurol*, 72(1), 41-52. doi:10.1002/ana.23557

- Morley J, & Tumosa N. (2002). Saint Louis University Mental Status Examination (SLUMS). *Aging Successfully*, 12(1), 4.
- Murray SA, & Sheikh A. (2008). Palliative care beyond cancer: Care for all at the end of life. *BMJ*, 336(7650), 958-959. doi:10.1136/bmj.39535.491238.94
- Nasreddine ZS, Phillips NA, Bedirian V, Charbonneau S, Whitehead V, Collin I, Cummings JL, & Chertkow H. (2005). The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment. *J Am Geriatr Soc*, 53(4), 695-699. doi:10.1111/j.1532-5415.2005.53221.x
- National Hospice Organization. (1996). Medical guidelines for determining prognosis in selected non-cancer diseases. *Hosp J*, 11(2), 47-63.
- Nguyen QA, & Paton C. (2008). The use of aromatherapy to treat behavioural problems in dementia. *Int J Geriatr Psychiatry*, 23(4), 337-346. doi:10.1002/gps.1886
- Nicholls D, Chang E, Johnson A, & Edenborough M. (2013). Touch, the essence of caring for people with end-stage dementia: a mental health perspective in Namaste Care. *Aging Ment Health*, 17(5), 571-578. doi:10.1080/13607863.2012.751581
- Olanzapine [package insert]. (2015). North Wales, PA: Teva Pharmaceuticals USA, Inc.
- Onega LL, Pierce TW, & Epperly L. (2016). Effect of bright light exposure on depression and agitation in older adults with dementia. *Issues Ment Health Nurs*, 37(9), 660-667. doi:10.1080/01612840.2016.1183736
- Petersen RC. (2011). Clinical practice. Mild cognitive impairment. *N Engl J Med*, 364(23), 2227-2234. doi:10.1056/NEJMcp0910237
- Reisberg B, Doody R, Stoffler A, Schmitt F, Ferris S, & Mobius HJ. (2003). Memantine in moderate-to-severe Alzheimer's disease. *N Engl J Med*, 348(14), 1333-1341. doi:10.1056/NEJMoa013128
- Reisberg B, Ferris SH, de Leon MJ, & Crook T. (1982). The Global Deterioration Scale for assessment of primary degenerative dementia. *Am J Psychiatry*, 139(9), 1136-1139. doi:10.1176/ajp.139.9.1136
- Reuben DB, Herr KA, Pacala JT, Pollock BG, Potter JF, & Semla TP. (2014). *Geriatrics at your fingertips*. 16th ed. Malden, MA: Blackwell.
- Reus VI, Fochtmann LJ, Eyler AE, Hilty DM, Horvitz-Lennon M, Jibson MD, Lopez OL, Mahoney J, Pasic J, Tan ZS, Wills CD, Rhoads R, & Yager J. (2016). The American Psychiatric Association Practice Guideline on the Use of Antipsychotics to Treat Agitation or Psychosis in Patients With Dementia. *American Journal of Psychiatry*, 173(5), 543-546. doi:10.1176/appi.ajp.2015.173501
- Risperidone [package insert]. (2016). Toronto, Ontario: Apotex Inc.
- Robinson L, Tang E, & Taylor JP. (2015). Dementia: timely diagnosis and early intervention. *BMJ*, 350, h3029. doi:10.1136/bmj.h3029
- Sachs GA, Shega JW, & Cox-Hayley D. (2004). Barriers to excellent end-of-life care for patients with dementia. *J Gen Intern Med*, 19(10), 1057-1063. doi:10.1111/j.1525-1497.2004.30329.x
- Sampson EL, Candy B, & Jones L. (2009). Enteral tube feeding for older people with advanced dementia. *Cochrane Database Syst Rev*(2), Cd007209. doi:10.1002/14651858.CD007209.pub2
- Schneider LS, Tariot PN, Dagerman KS, Davis SM, Hsiao JK, Ismail MS, Lebowitz BD, Lyketsos CG, Ryan JM, Stroup TS, Sultzer DL, Weintraub D, & Lieberman

JA. (2006). Effectiveness of atypical antipsychotic drugs in patients with Alzheimer's disease. *N Engl J Med*, 355(15), 1525-1538. doi:10.1056/NEJMoa061240

Schulz R, Mendelsohn AB, Haley WE, Mahoney D, Allen RS, Zhang S, Thompson L, & Belle SH. (2003). End-of-life care and the effects of bereavement on family caregivers of persons with dementia. *N Engl J Med*, 349(20), 1936-1942. doi:10.1056/NEJMsa035373

Sclan SG, & Reisberg B. (1992). Functional assessment staging (FAST) in Alzheimer's disease: reliability, validity, and ordinality. *Int Psychogeriatr*, 4 Suppl 1, 55-69.

Sellers DM. (2006). The evaluation of an animal assisted therapy intervention for elders with dementia in long-term care. *Activities, Adaptation & Aging*, 30(1), 61-77. doi:10.1300/J016v30n01\_04

Semla T, Beizer J, & Higbee M. (2007). *Geriatric Dosage Handbook* (12th ed.). Hudson, OH: Lexi-Comp, Inc.

Shega J, & Levine S. (2012). Caring for patients with chronic illnesses: dementia, COPD, and CHF. In P. J. Storey (Ed.), *Unipac 9: a resource for hospice and palliative care professionals* (pp. 1-33). Glenview, IL: American Academy of Hospice and Palliative Medicine.

Simmons BB, Hartmann B, & DeJoseph D. (2011). Evaluation of suspected dementia. *Am Fam Physician*, 84(8), 895-902.

Slaughter SE, Eliasziw M, Morgan D, & Drummond N. (2011). Incidence and predictors of eating disability among nursing home residents with middle-stage dementia. *Clin Nutr*, 30(2), 172-177. doi:10.1016/j.clnu.2010.09.001

Street JS, Clark WS, Gannon KS, Cummings JL, Bymaster FP, Tamura RN, Mitan SJ, Kadam DL, Sanger TM, Feldman PD, Tollefson GD, & Breier A. (2000). Olanzapine treatment of psychotic and behavioral symptoms in patients with Alzheimer disease in nursing care facilities: a double-blind, randomized, placebo-controlled trial. The HGEU Study Group. *Arch Gen Psychiatry*, 57(10), 968-976.

Streim JE, Porsteinsson AP, Breder CD, Swanink R, Marcus R, McQuade R, & Carson WH. (2008). A randomized, double-blind, placebo-controlled study of aripiprazole for the treatment of psychosis in nursing home patients with Alzheimer disease. *Am J Geriatr Psychiatry*, 16(7), 537-550. doi:10.1097/JGP.0b013e318165db77

Tariot PN, Schneider L, Katz IR, Mintzer JE, Street J, Copenhaver M, & Williams-Hughes C. (2006). Quetiapine treatment of psychosis associated with dementia: a double-blind, randomized, placebo-controlled clinical trial. *Am J Geriatr Psychiatry*, 14(9), 767-776. doi:10.1097/01.jgp.0000196628.12010.35

Teno JM, Gozalo PL, Bynum JP, Leland NE, Miller SC, Morden NE, Scupp T, Goodman DC, & Mor V. (2013). Change in end-of-life care for Medicare beneficiaries: site of death, place of care, and health care transitions in 2000, 2005, and 2009. *JAMA*, 309(5), 470-477. doi:10.1001/jama.2012.207624

Tegeler, M., Miller, E., Thomas, A., & Pemberton, A. (2015). OPTIMISTIC Interprofessional Dementia Training: Strategies to Manage Challenging Behaviors. *Journal of the American Medical Directors Association*, 16(3), B14.

Threlfall A, Barton C, & Yaffe K. (2013). Dementia. In S. Durso & G. Sullivan (Eds.), *Geriatrics Review Syllabus: A Core Curriculum in Geriatric Medicine* (8th ed., pp. 284-295). New York, NY: American Geriatrics Society.

Whear R, Coon JT, Bethel A, Abbott R, Stein K, & Garside R. (2014). What is the impact of using outdoor spaces such as gardens on the physical and mental well-being of those with dementia? A systematic review of quantitative and qualitative evidence. *J Am Med Dir Assoc*, 15(10), 697-705. doi:10.1016/j.jamda.2014.05.013

Winblad B, Kilander L, Eriksson S, Minthon L, Batsman S, Wetterholm AL, Jansson-Blixt C, & Haglund A. (2006). Donepezil in patients with severe Alzheimer's disease: double-blind, parallel-group, placebo-controlled study. *Lancet*, 367(9516), 1057-1065. doi:10.1016/s0140-6736(06)68350-5

Zanni GR, & Wick JY. (2007). Differentiating dementias in long-term care patients. *Consult Pharm*, 22(1), 14-16, 18, 21-18.